arXiv:1512.07479v1 [math.GN] 23 Dec 2015

Metrizable compacta in the space of continuous functions with the topology of pointwise convergence

V.V. Mykhaylyuk [1]

Chernivtsi National University, Department of Mathematical Analysis, Kotsjubyns'koho 2, Chernivtsi 58012, Ukraine
e-mail: mathan@ukr.net

Abstract

We prove that every point-finite family of nonempty functionally open sets in a topological space X has the cardinality at most an infinite cardinal κ if and only if $w(X) \leq \kappa$ for every Valdivia compact space $Y \subseteq C_p(X)$. Correspondingly a Valdivia compact space Y has the weight at most an infinite cardinal κ if and only if every point-finite family of nonempty open sets in $C_p(Y)$ has the cardinality at most κ, that is $p(C_p(Y)) \leq \kappa$. Besides, it was proved that $w(Y) = p(C_p(Y))$ for every linearly ordered compact Y. In particular, a Valdivia compact space or linearly ordered compact space Y is metrizable if and only if $p(C_p(Y)) = \aleph_0$. This gives answer to a question of O. Okunev and V. Tkachuk.

1 Introduction

During the study of the Baire classification of separately continuous functions $f : X \times Y \to \mathbb{R}$, when Y satisfies compactness type conditions, it was established that spaces X which satisfy a condition of a countable chain type, play an important role (see [1,2,3]). The Baire classification of these mappings was obtained replacing of Y with its continuous metrizable image in the space $C_p(X)$ of all continuous functions on X with the pointwise convergence topology.

This shows that it is natural to try to find necessary and sufficient conditions on a topological space X for metrizability of any compact subspace of $C_p(X)$

[1] *Key words and phrases:* Valdivia compact, metrizable compact, separately continuous function, point-finite cellularity, linearly ordered compact.
2000 Mathematics Subject Classification: Primary 54E45, 54C35; Secondary 54A25, 54D30

or any compact subspace of $C_p(X)$ which belongs to a given class such as Eberlein compact spaces, Corson compact spaces or Valdivia compact spaces.

A compact space X which is homeomorphic to a weakly compact subset of a Banach space is called *an Eberlein compact space*. A compact space X is called *a Corson compact space* if it is homeomorphic to a compact space $Z \subseteq \mathbb{R}^T$ such that $|\operatorname{supp} z| \leq \aleph_0$ for any $z \in Z$. By $\operatorname{supp} z$ we denote the support $\{t \in T : z(t) \neq 0\}$ of a function $z : T \to \mathbb{R}$. A compact space X is called *a Valdivia compact space* if it is homeomorphic to a compact space $Z \subseteq \mathbb{R}^T$ such that the set $\{z \in Z : |\operatorname{supp} z| \leq \aleph_0\}$ is dense in Z.

Analogous questions were investigated in [4,5,6,7],where connections between cardinal properties of a completely regular space X and the space $C_p(X)$ were studied. The point-finite cellularity $p(X) = \sup\{|\tau| : \tau$ is a point-finite family of nonempty open sets in $X\}$ of a topological space X, which was introduced in [4], is important in these papers. In particular, the following questions were posed in [7, questions 5.2, 5.4]:

Question 1.1 *Let X be a Souslin continuum, i.e. a non-separable linearly ordered perfectly normal compact space. It is true that $p(C_p(X)) = \aleph_0$?*

Question 1.2 *Let X be a Corson compact space such that $p(C_p(X)) = \aleph_0$. Must X be metrizable?*

Question 1.2 was answered in [8]. It is proved in [8] that the weight of a Corson compact space X is equal to the point-finite cellularity of $C_p(X)$.

In this paper we study necessary and sufficient conditions on a topological space X for metrizability of any Valdivia compact space $Y \subseteq C_p(X)$. Using a technique of a dependence of mappings upon a certain number of coordinates we prove that any point-finite family of functionally open sets in a topological space X has the cardinality at most an infinite cardinal κ if and only if each Valdivia compact space $Y \subseteq C_p(X)$ has the topological weight at most κ. In particular, $p(X) = \aleph_0$ for a completely regular space X if and only if each Valdivia compact space $Y \subseteq C_p(X)$ is metrizable. Passing to a dual problem on study of connections between properties of a compact space Y and the space $C_p(Y)$, we obtain that a Valdivia compact space Y has the weight at most an infinite cardinal κ if and only if $p(C_p(Y)) \leq \kappa$. It implies immediately the following fact: a Valdivia compact space Y is metrizable if and only if $p(C_p(Y)) = \aleph_0$. Besides, it was proved that $w(Y) = p(C_p(Y))$ for every linearly ordered compact space Y (this yields a negative answer to Question 5.2 from [7]).

2 A dependence of mappings upon a certain number of coordinates

In this section we prove a theorem which will be a technical tool in the proof of the main result and based on a notion of a dependence of mappings upon a certain number of coordinates.

Let κ be an infinite cardinal number, and let Z, T be sets, $Y \subseteq \mathbb{R}^T$ and $f : Y \to Z$. We say that f *is concentrated on a set* $S \subseteq T$ if for any $y', y'' \in Y$ the condition $y'|_S = y''|_S$ implies $f(y') = f(y'')$. Besides, if $|S| \leq \kappa$ then f *depends upon* κ *coordinates.*

Let X be a set and $g : X \times Y \to Z$. Then g *is concentrated on a set* $S \subseteq T$ *in the second variable* if $g(x, y') = g(x, y'')$ for any $x \in X$ and $y', y'' \in Y$ with $y'|_S = y''|_S$; and g *depends upon* κ *coordinates in the second variable* if $|S| \leq \kappa$ for some S.

A subset A of a topological space X is called *functionally open* if there exists a continuous function $f : X \to [0, 1]$ such that $A = f^{-1}((0, 1])$.

The following result was obtained in [9] for $\kappa = \aleph_0$.

Theorem 2.1 *Let X be a topological space in which every point-finite family of nonempty functionally open sets has the cardinality at most an infinite cardinal number κ, let $Y \subseteq \mathbb{R}^T$ be a compact space, $B = \{y \in Y : |\operatorname{supp} y| \leq \kappa\}$ and $Y = \overline{B}$. Then every separately continuous function $f : X \times Y \to \mathbb{R}$ depends upon κ coordinates in the second variable.*

Proof. First we prove that for every $\varepsilon > 0$ there exists a set $R_\varepsilon \subseteq T$ such that $|R_\varepsilon| \leq \kappa$ and for any $b', b'' \in B$ the condition $b'|_{R_\varepsilon} = b''|_{R_\varepsilon}$ implies $|f(x, b') - f(x, b'')| \leq \varepsilon$ for every $x \in X$.

Suppose, contrary to our claim, that there exists an $\varepsilon > 0$ such that for every set $S \subseteq T$ with $|S| \leq \kappa$ there exist an $x \in X$ and $b', b'' \in B$ such that $b'|_S = b''|_S$ and $|f(x, b') - f(x, b'')| > \varepsilon$.

Denote by $T(\kappa^+)$ the first ordinal of cardinality κ^+. Using the transfinite induction we construct a system $\{S_\alpha : \alpha < T(\kappa^+)\}$ of sets $S_\alpha \subseteq T$ and sets $\{b_\alpha : \alpha < T(\kappa^+)\}$, $\{c_\alpha : \alpha < T(\kappa^+)\}$ and $\{x_\alpha : \alpha < T(\kappa^+)\}$ of points $b_\alpha, c_\alpha \in B$ and $x_\alpha \in X$ such that

(a) $|S_\alpha| \leq \kappa$ for any $\alpha < T(\kappa^+)$;

(b) $b_\alpha|_{S_\alpha} = c_\alpha|_{S_\alpha}$ for any $\alpha < T(\kappa^+)$;

(c) $S_\alpha \subseteq S_\beta$ for any $\alpha < \beta < T(\kappa^+)$;

(d) $\operatorname{supp} b_\alpha \subseteq S_{\alpha+1}$, $\operatorname{supp} c_\alpha \subseteq S_{\alpha+1}$ for any $\alpha < T(\kappa^+)$;

(e) $|f(x_\alpha, b_\alpha) - f(x_\alpha, c_\alpha)| > \varepsilon$ for any $\alpha < T(\kappa^+)$.

Fix a set $S_1 \subseteq T$ with $|S_1| \leq \kappa$. By the assumption, there exist points $x_1 \in X$ and $b_1, c_1 \in B$ such that $b_1|_{S_1} = c_1|_{S_1}$ and $|f(x_1, b_1) - f(x_1, c_1)| > \varepsilon$. Put $S_2 = S_1 \cup \operatorname{supp} b_1 \cup \operatorname{supp} c_1$. Clearly $|S_2| \leq \kappa$. Choose points $x_2 \in X$ and $b_2, c_2 \in B$ such that $b_2|_{S_2} = c_2|_{S_2}$ and $|f(x_2, b_2) - f(x_2, c_2)| > \varepsilon$.

Suppose that for some $\beta < T(\kappa^+)$ the sets $\{S_\alpha : \alpha < \beta\}$, $\{b_\alpha : \alpha < \beta\}$, $\{c_\alpha : \alpha < \beta\}$ and $\{x_\alpha : \alpha < \beta\}$ are constructed. Put $S_\beta = \bigcup\limits_{\alpha<\beta} (S_\alpha \cup \operatorname{supp} b_\alpha \cup \operatorname{supp} c_\alpha)$. Since for $\alpha < \beta$ all the sets S_α, $\operatorname{supp} b_\alpha$ and $\operatorname{supp} c_\alpha$ have cardinality at most κ, we have the inequality $|S_\beta| \leq \kappa$. By the assumption, there exist points x_β and $b_\beta, c_\beta \in B$ such that $b_\beta|_{S_\beta} = c_\beta|_{S_\beta}$ and $|f(x_\beta, b_\beta) - f(x_\beta, c_\beta)| > \varepsilon$.

The continuity of f in variable x and condition (e) imply that for every $\alpha < T(\kappa^+)$ there exists an functionally open neighborhood U_α of x_α in X such that $|f(x, b_\alpha) - f(x, c_\alpha)| > \varepsilon$ for any $x \in U_\alpha$. According to the theorem condition, the family $(U_\alpha : \alpha < \omega)$ is not point-finite. Thus there exist a point $x_0 \in X$ and a strictly increasing sequence $(\alpha_n)_{n=1}^\infty$ of ordinals $\alpha_n < T(\kappa^+)$ such that $|f(x_0, b_{\alpha_n}) - f(x_0, c_{\alpha_n})| > \varepsilon$ for any $n \in \mathbb{N}$.

Put $T_n = S_{\alpha_n}$, $v_n = b_{\alpha_n}$ and $w_n = c_{\alpha_n}$ by $n \in \mathbb{N}$. Using the compactness of Y and the continuity of $f^{x_0} : Y \to \mathbb{R}$, $f^{x_0}(y) = f(x_0, y)$, choose a finite set $T_0 \subseteq T$ such that $|f(x_0, y') - f(x_0, y'')| < \varepsilon$ for any $y', y'' \in Y$ with $y'|_{T_0} = y''|_{T_0}$. It follows from $|f(x_0, v_n) - f(x_0, w_n)| > \varepsilon$ that $v_n|_{T_0} \neq w_n|_{T_0}$. But $v_n|_{T_n} = w_n|_{T_n}$ according to (b) and $v_n|_{T \setminus T_{n+1}} = w_n|_{T \setminus T_{n+1}}$ according to (c) and (d). Thus $T_0 \cap (T_{n+1} \setminus T_n) \neq \emptyset$ for any $n \in \mathbb{N}$. Since the sequence $(T_n)_{n=1}^\infty$ is increasing, T_0 is infinite, which is impossible. Hence the existence of R_ε is proved.

Put $R_0 = \bigcup\limits_{n=1}^{\infty} R_{\frac{1}{n}}$. Clearly $f(x, b') = f(x, b'')$ for any $x \in X$, $b', b'' \in B$ with $b'|_{R_0} = b''|_{R_0}$. Fix points $x \in X$, $y', y'' \in Y$ such that $y'|_{R_0} = y''|_{R_0}$. Since a function $f^x : Y \to \mathbb{R}$, $f^x(y) = f(x, y)$ is continuous on the compact space $Y \subseteq \mathbb{R}^T$, there exists an at most countable set $T_0 \subseteq T$ such that $f(x, y_1) = f(x, y_2)$ for any $y_1, y_2 \in Y$ with $y_1|_{T_0} = y_2|_{T_0}$. Consider a continuous mapping $\varphi : Y \to \mathbb{R}^{T_0 \cup R_0}$, $\varphi(y) = y|_{T_0 \cup R_0}$. Since $Y = \overline{B}$, we have the inequality $\varphi(Y) = \overline{\varphi(B)}$. Note that B is an κ-compact set, that is, the closure $\overline{A}$ in B of any set $A \subseteq B$ with $|A| \leq \kappa$ is a compact set. Therefore $\varphi(B)$ is an κ-compact set. It follows from $|T_0 \cup R_0| \leq \kappa$ that $w(\varphi(Y)) \leq \kappa$. The density of $\varphi(B)$ in $\varphi(Y)$ implies the existence of a dense in $\varphi(Y)$ set $C \subseteq \varphi(B)$ such that $|C| \leq \kappa$. Then $\varphi(Y) \subseteq \overline{C} \subseteq \varphi(B)$. Thus $\varphi(Y) = \varphi(B)$. Therefore there exist points $b', b'' \in B$ such that $b'|_{T_0 \cup R_0} = y'|_{T_0 \cup R_0}$ and $b''|_{T_0 \cup R_0} = y''|_{T_0 \cup R_0}$. Since $b'|_{T_0} = y'|_{T_0}$, $b''|_{T_0} = y''|_{T_0}$ and $b'|_{R_0} = b''|_{R_0}$, we have $f(x, y') = f(x, b')$, $f(x, b'') = f(x, y'')$ and $f(x, b') = f(x, b'')$. Hence $f(x, y') = f(x, y'')$. Thus f is

concentrated on R_0 and therefore f depends upon κ coordinates in the second variable. $\diamondsuit$

3 Main results

Theorem 3.1 *Let X be a topological space and let κ be an infinite cardinal number. Then the following conditions are equivalent:*

(i) every point-finite family of nonempty functionally open in X sets has the cardinality at most κ;

(ii) $w(Y) \leq \kappa$ for any Valdivia compact space $Y \subseteq C_p(X)$;

(iii) $w(Y) \leq \kappa$ for any Eberlein compact space $Y \subseteq C_p(X)$.

Proof. $(i) \Longrightarrow (ii)$. By the definition, a Valdivia compact space $Y \subseteq C_p(X)$ is homeomorphic to a compact space $Z \subseteq \mathbb{R}^T$ such that the set $\{z \in Z : |\operatorname{supp} z| \leq \aleph_0\}$ is dense in Z. Fix a homeomorphism $\varphi : Z \to Y$. For every $x \in X$ and $z \in Z$ we put $f(x, z) = \varphi(z)(x)$. Clearly, f is a separately continuous function on $X \times Z$. By Theorem 2.1, f depends upon κ coordinates in the second variable. This implies that the map $\psi : Z \to \mathbb{R}^S$, $\psi(z) = z|_S$, is an injection and hence homeomorphism onto $\tilde{Z} = \psi(Z)$. Therefore $w(Z) = w(\tilde{Z}) \leq \kappa$.

Since every Eberlein compact space is a Valdivia compact space, the implication $(ii) \Longrightarrow (iii)$ is obvious.

The implication $(iii) \Longrightarrow (i)$ was proved in [5, Theorem 1].

Corollary 3.2 *Let X be a completely regular space. Then the following conditions are equivalent:*

(i) $p(X) = \aleph_0$;

(ii) each Valdivia compact space $Y \subseteq C_p(X)$ is metrizable.

A topological space X is called *pseudocompact* if every continuous function $f : X \to \mathbb{R}$ is bounded; *countably compact* if a finite subcover can be chosen from any countable open cover of X. A topological space X is called *σ-pseudocompact* if it is the union of a sequence of pseudocompact subspaces of X.

Corollary 3.3 *Let X be a completely regular space with $p(X) = \aleph_0$ which consists a σ-pseudocompact dense subspace. Then every countably compact*

space $Y \subseteq C_p(X)$ is metrizable.

Proof. By [10, Theorem III.4.23] every countably compact space $Y \subseteq C_p(X)$ is an Eberlein compact space. It remains to use Corollary 3.2. $\diamondsuit$

By $\mathcal{K}(X)$ we denote the system of all compact subsets of a topological space X. A topological space X is called *$\mathcal{K}$-countably-determined* if there exist a subset Y in the topological space $\mathbb{N}^{\mathbb{N}}$ and a set-valued mapping $F : Y \to \mathcal{K}(X)$ such that for every open in X set U the set $\{y \in Y : F(y) \subseteq U\}$ is open in Y and $X = \bigcup\limits_{y \in Y} F(y)$.

Corollary 3.4 *Let X be a completely regular space with $p(X) = \aleph_0$ which contains a $\mathcal{K}$-countably-determined dense subspace. Then every compact space $Y \subseteq C_p(X)$ is metrizable.*

Proof. Fix a dense in X $\mathcal{K}$-countably-determined subspace $\tilde{X}$ and a compact space $Y \subseteq C_p(X)$. Consider the continuous identity mapping $\varphi : C_p(X) \to C_p(\tilde{X})$. Put $\tilde{Y} = \varphi(Y)$. It follows from density $\tilde{X}$ in X that $\varphi|_Y$ is a bijection and by [11, Theorem 3.1.13] φ is a homeomorphism. By [12, Theorem 3.7] $\tilde{Y}$ is a Corson compact space. Thus Y is a Corson compact space and by Corollary 3.2 it is metrizable. $\diamondsuit$

Let D be a discrete space of cardinality $\kappa > \aleph_0$. By L_κ we will denote an one-point Lindelöffication of D, that is $L_\kappa = D \bigcup \{\infty\}$ and all neighborhoods of ∞ in L_κ are the complements in L_κ to all at most countable subsets of D, besides the discrete topology on D coincides with the topology on D induced by L_κ. A topological space X is called *primary Lindelöff* if X is a continuous image of a closed subset of the topological product ${L_\kappa}^{\mathbb{N}}$ for some $\kappa > \aleph_0$. Note that by [10, Corollary IV.3.17] for a primary Lindelöff space X every compact space $Y \subseteq C_p(X)$ is a Corson compact space.

The following result can be obtained analogously.

Corollary 3.5 *Let X be a completely regular space with $p(X) = \aleph_0$ which contains a primary Lindelöff dense subspace. Then every compact space $Y \subseteq C_p(X)$ is metrizable.*

Recall that a set F of functions defined on a set X *separates points on X* if for arbitrary distinct points $x', x'' \in X$ there exists a function $f \in F$ such that $f(x') \neq f(x'')$.

Theorem 3.6 *Let Y be a Valdivia compact space and let κ be an infinite cardinal number. Then the following conditions are equivalent:*

(i) $w(Y) \leq \kappa$;

(ii) $p(C_p(Y)) \leq \kappa$;

(iii) there exists a space $X \subseteq C_p(Y)$ *with* $p(X) \leq \kappa$ *which separates points on* Y.

Proof. $(i) \Longrightarrow (ii)$. Suppose that $w(Y) \leq \kappa$. If Y is infinite then $d(C_p(Y)) \leq w(Y) \leq \kappa$. If Y is finite then $d(C_p(Y)) = \aleph_0 \leq \kappa$ anyway. Thus $d(C_p(Y)) \leq \kappa$. Therefore $p(C_p(Y)) \leq \kappa$.

The implication $(ii) \Longrightarrow (iii)$ is obvious.

$(iii) \Longrightarrow (i)$. Fix a subspace $X \subseteq C_p(Y)$ with $p(X) \leq \kappa$ which separates points on Y. Consider a continuous mapping $\varphi : Y \to C_p(X)$, $\varphi(y)(x) = x(y)$. Since X separates points on Y, the map $\varphi : Y \to \varphi(Y)$ is a bijection and by [11, Theorem 3.1.13] φ is a homeomorphism. Thus $\varphi(Y)$ is an Valdivia compact space and $w(Y) \leq \kappa$ by Theorem 3.1. $\diamondsuit$

Note that Theorem 3.6 is not true for an arbitrary compact space Y. Indeed, $2^{\aleph_0} = w(Y) > p(C_p(Y)) = \aleph_0$ for $Y = \beta\mathbb{N}$ (see [6], proof of Theorem 2.11).

Corollary 3.7 *Any Valdivia compact space* Y *is metrizable if and only if* $p(C_p(Y)) = \aleph_0$.

4 Linearly ordered compacta

Let $(X, <)$ be a linearly ordered space. For every $x, y \in X$, $x < y$, we put $(x, y) = \{z \in X : x < z < y\}$, $[x, y) = \{z \in X : x \leq z < y\}$, $(x, y] = \{z \in X : x < z \leq y\}$ and $[x, y] = \{z \in X : x \leq z \leq y\}$. Elements $x, y \in X$, $x < y$ such that $(x, y) = \emptyset$ are called *neighbor points*.

Proposition 4.1 *Let* $(X, <)$ *be a linearly ordered countably compact space and let* $A = \{(x, y) \in X^2 : x < y \text{ and } (x, y) = \emptyset\}$. *Then* $c(X) \leq p(C_p(X))$ *and* $|A| \leq p(C_p(X))$.

Proof. Let $B = \{(x_i, y_i) \in X^2 : i \in I\}$ be a set which satisfies the following conditions:

a) $x_i < y_i$ for every $i \in I$;

b) $y_i \leq x_j$ or $y_j \leq x_i$ for every distinct $i, j \in I$.

Put $U_i = \{f \in C_p(X) : f(x_i) < 0 \text{ and } f(y_i) > 1\}$ for each $i \in I$. We show that $(U_i : i \in I)$ is a point-finite family in $C_p(X)$. Suppose that $f \in C_p(X)$ such that a set $J = \{i \in I : f \in U_i\}$ is infinite. Then in an infinite set $\{x_i : i \in J\}$

there exists a strictly increasing or decreasing sequence $(x_{i_n})_{n=1}^{\infty}$, $i_n \in J$. Since X is a countably compact space, there exists $z \in X$ such that $z = \lim\limits_{n\to\infty} x_{i_n} = \lim\limits_{n\to\infty} y_{i_n}$. Therefore $f(z) = \lim\limits_{n\to\infty} f(x_{i_n}) \leq 0$ and $f(z) = \lim\limits_{n\to\infty} f(y_{i_n}) \geq 1$, which is impossible.

Thus $(U_i : i \in I)$ is a point-finite family and $|I| \leq p(C_p(X))$. Since any chain generates a set B with $(x_i, y_i) \neq \emptyset$ for every $i \in I$, and A generates a set B with $(x_i, y_i) = \emptyset$ for every $i \in I$, we have the inequality $c(X) \leq p(C_p(X))$ and $|A| \leq p(C_p(X))$. $\diamondsuit$

The following assertion seems to be known.

Proposition 4.2 *Let $(X, <)$ be a linearly ordered space. Then $d(X) \leq c(X^2)$.*

Proof. Let A be a set of all isolated points of X and let $(W_i : i \in I)$ be a maximal family of nonempty sets $W_i = (a_i, b_i) \times (c_i, d_i)$ in X^2 such that $b_i \leq c_i$ or $d_i \leq a_i$ for each $i \in I$. Clearly $|A| \leq c(X^2)$ and $|I| \leq c(X^2)$. Put $B = A \cup \{a_i, b_i, c_i, d_i : i \in I\}$. Suppose that $U = X \setminus \overline{B} \neq \emptyset$.

Choose a nonempty set $(x, y) \subseteq U$. Since $U \cap A = \emptyset$ then there exist $a, b, c, d \in (x, y)$ such that $a < b \leq c < d$ and sets (a, b) and (c, d) are nonempty. Then an open in X^2 nonempty set $W = (a, b) \times (c, d)$ such that $W \cap W_i = \emptyset$ for every $i \in I$. This contradicts the maximality of $(W_i : i \in I)$. $\diamondsuit$

Theorem 4.3 *Let $(X, <)$ be an infinite linearly ordered compact. Then $w(X) = p(C_p(X))$.*

Proof. The inequality $p(C_p(X)) \leq w(X)$ can be proved like in the proof of Theorem 3.6.

Let $p(C_p(X)) = \kappa$. Then a set $\tilde{A} = \{(x, y) \in X^2 : x < y \ \textit{and} \ (x, y) = \emptyset\}$ has the cardinality at most κ by Proposition 4.1. Therefore a set $A = \{x, y \in X : (x, y) \in \tilde{A}\}$ has the cardinality at most κ. Note that for arbitrary dense in X set B the system $\{(x, y) : x < y,\ x, y \in A \cup B\} \bigcup \{[a, x) : x \in A \cup B\} \bigcup \{(x, b] : x \in A \cup B\}$, where $a = \min X$ and $b = \max X$, is a base of open in X sets. Therefore it is sufficient to prove that $d(X) \leq \kappa$.

Consider a system $\mathcal{V}$ of all open in X nonempty set V with $d(V) \leq \kappa$. Choose a maximal disjoint system $\mathcal{U} \subseteq \mathcal{V}$. Proposition 4.1 implies $|\mathcal{U}| \leq \kappa$. For every $U \in \mathcal{U}$ we choose a set $B_U \subseteq U$ such that $|B_U| \leq \kappa$ and $U \subseteq \overline{B_U}$. Put $B = A \bigcup \bigcup\limits_{U \in \mathcal{U}} B_U$ and $Y = \overline{B}$. Then $|B| \leq \kappa^2 = \kappa$.

Suppose that $X \setminus Y \neq \emptyset$. Choose $x_0, y_0 \in X \setminus Y$ such that $x_0 < y_0$ and $X_0 = [x_0, y_0] \subseteq X \setminus Y$. We show that $c(X_0^2) \leq \kappa$.

Let $(W_i : i \in I)$ be a disjoint family of open sets in X_0^2. Suppose that $|I| > \kappa$. For every $i \in I$ choose pair-wise distinct points $a_i, b_i, c_i, d_i \in X_0$ such that $a_i < b_i$, $c_i < d_i$ and $(a_i, b_i) \times (c_i, d_i) \subseteq W_i$ (it is possible since X_0 has no neighbor point). Put $U_i = \{f \in C_p(X_0) : f(a_i) < 0, f(b_i) > 1, f(c_i) < 0, f(d_i) > 1\}$ for every $i \in I$. Note that $p(C_p(X_0)) \leq \kappa$, thus $(U_i : i \in I)$ is not point-finite in $C_p(X_0)$. There exist a function $f_0 \in C_p(X_0)$ and an infinite set $J \subseteq I$ such that $f_0 \in U_i$ for every $i \in J$. Since f_0 is continuous on linearly ordered compact space X_0 which has no neighbor point, there exists a finite set $K \subseteq X_0$ such that for every $x, y \in X_0$ the inequality $|f_0(x) - f_0(y)| > 1$ implies $(x, y) \cap K \neq \emptyset$. Then $(a_i, b_i) \cap K \neq \emptyset$, $(c_i, d_i) \cap K \neq \emptyset$ therefore $W_i \cap K^2 \neq \emptyset$ for every $i \in J$ which contradicts the disjointness of $(W_i : i \in I)$. Hence $|I| \leq \kappa$.

Thus $c(X_0^2) \leq \kappa$ and according to Proposition 4.2 we obtain that $d(X_0) \leq \kappa$, which contradicts the maximality of the system $\mathcal{U}$. Therefore $X \setminus Y = \emptyset$, $d(X) \leq \kappa$ and $w(X) \leq \kappa$. $\diamondsuit$

The following corollary gives a negative answer to Question 5.2 from [7].

Corollary 4.4 *A linearly ordered compact X is metrizable if and only if $p(C_p(X)) = \aleph_0$.*

References

[1] W. Moran *Separate continuity and support of measures*, J. London. Math. Soc. **44** (1969), 320-324.

[2] G. Vera *Baire measurability of separately continuous functions*, Quart. J. Math. Oxford. **39, 153** (1988), 109-116.

[3] V.V. Mykhaylyuk, O.V. Sobchuk *Baire classification of separately continuous functions and dependence upon countable number of coordinates*, Nauk. Visn. Chern. Univ. **Vyp. 191-192**, Matem., Chernivtsi: Ruta (2004), 116-118 (in Ukrainian).

[4] V.V. Tkachuk *Cardinal invariants of the Suslin number type*, Dokl. Akad. Nauk SSSR. **270, 4** (1983), 795-798 (in Russian).

[5] V.V. Tkachuk *A supertopological cardinal invariants*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. **4** (1984), 26-29(in Russian).

[6] A.V. Arhangel'skii, V.V. Tkachuk V.V. *Calibers and point-finite cellularity of the space $C_p(X)$ and some questions of S.Gul'ko and M.Husek*, Topology Appl. **23, 1** (1986), 65-73.

[7] O.G. Okunev, V.V. Tkachuk *Lindelöf Σ-property in $C_p(X)$ and $p(C_p(X)) = \omega$ do not imply countable network weight in X*, Acta Math. Hungar. **90, 1-2** (2001), 119-132.

[8] N.D. Kalamidas, G.D. Spiliopoulos *Compact sets in $C_p(X)$ and calibers*, Can. Math. Bull. **35, 4** (1992), 497-502.

[9] V.V. Mykhaylyuk *One-point discontinuity set of separately continuous functions on the product of two compact spaces*, Ukr. Mat. Zhurn. **57, 1** (2005), 94-101 (in Ukrainian).

[10] A.V. Arhangelskii *Topological spaces of functions*, .: Izd-vo Moskovskogo Univ. (1989), 222 s.(in Russian).

[11] R. Engelking *General topology*, M.: Mir (1986), 752 s. (in Russian).

[12] M. Talagrand *Espaces de Banach faiblement $\mathcal{K}$-analytiques*, Ann. of Math. **110** (1979), 407-438.